



## PHYS 3317: Phonons A

Hand in at beginning of next lecture (only do handout A, B, or C, not all three)

### Problem 1. Kinetic Energy

1.1. We defined creation and annihilation operators via

$$x_j = \frac{d}{\sqrt{2}} \left( [(a_j + a_j^\dagger) + \alpha(a_{j+1} + a_{j+1}^\dagger + a_{j-1} + a_{j-1}^\dagger)] \right) \quad (1)$$

$$p_j = \frac{\hbar}{\sqrt{2}di} \left( [(a_j - a_j^\dagger) - \alpha(a_{j+1} - a_{j+1}^\dagger + a_{j-1} - a_{j-1}^\dagger)] \right). \quad (2)$$

where  $\alpha$  is small. To leading order in  $\alpha$ , these obey

$$[a_i, a_j] = 0 \quad (3)$$

$$[a_i, a_j^\dagger] = \delta_{ij} \quad (4)$$

where  $\delta_{ij}$  is the Kronecker delta, equal to zero if  $i \neq j$  and 1 if  $i = j$ .

Write the kinetic energy in terms of these creation and annihilation operators. Neglect terms which are of quadratic or higher order in  $\alpha$ .

$$K = \sum_j \frac{p_j^2}{2m} \quad (5)$$

Rewrite your expression in the form:

$$K = \sum_j A_K (a_j^\dagger a_j + a_j a_j^\dagger) + B_K (a_{j+1}^\dagger a_j + a_j^\dagger a_{j+1}) + C_K (a_j a_j + a_j^\dagger a_j^\dagger) + D_K (a_{j+1} a_j + a_j^\dagger a_{j+1}^\dagger) \quad (6)$$

Note, to put it in this form, you will need to use the identity:

$$\sum_j c_{j-1} d_j = \sum_j c_j d_{j+1}. \quad (7)$$

